

A NEW *CLATHRIA* (PORIFERA: DEMOSPONGIAE: MICROCIONIDAE) FROM THE WESTERN INDIAN OCEAN

JOHN N.A. HOOPER, MICHELLE KELLY AND JOHN A. KENNEDY

Hooper, J.N.A., Kelly, M. & Kennedy, J.A. 2000 06 30: A new *Clathria* (Porifera: Demospongiae: Microcionidae) from the Western Indian Ocean. *Memoirs of the Queensland Museum* 42(2): 427-444. Brisbane. ISSN 0079-8835.

A new species of microcionid marine sponge *Clathria (Microciona) richmondi* sp. nov. is described from Zanzibar, Tanzania, and is highly unusual in having extremely large accolada toxas that form dragmata and skeletal tracts within the choanosomal and ectosomal skeletons, rudimentary spination on echinating acanthostyles, a live blue colouration and a prominent sub-surface aquiferous system with radiate arrangement around oscules. The new species is compared with the other 64 species of *Clathria* described from the Western Indian Ocean, Southeast Africa and Arabian Gulf-Red Sea provinces, and other species known to have toxodragmata. □ *Porifera, Demospongiae, Poecilosclerida, Microcionidae, new species, taxonomy, Zanzibar, Western Indian Ocean.*

John N.A. Hooper (email: JohnH@qm.qld.gov.au) and John A. Kennedy, Queensland Museum, PO Box 3300, South Brisbane 4101, Australia; Michelle Kelly, Marine Ecology and Aquaculture, National Institute of Water and Atmospheric Research (NIWA), Private Bag 109-695, Newmarket, Auckland, New Zealand; 14 December 1999.

The extensive literature on sponges of the Western Indian Ocean is evidence of a rich and diverse fauna (Table 1). Even though this inventory already contains 'several hundred' species (Kelly, 1997), it is undoubtedly far from complete. For example, Van Soest (1994b) collected 240 species from the Seychelles and Amirante islands whereas only 167 species were known previously from the region.

Within the Western Indian Ocean (i.e. extending along the continental shelf from Natal to Somalia and including the islands from Madagascar to the Seychelles; Richmond, 1997), Southeast African regions (Natal to the Cape) and Arabian Gulf-Red Sea provinces 74 species of Microcionidae have been reported (Hooper, 1996a), of which 64 belong to *Clathria*, 10 still unnamed (Table 2). Of these, 44 (or 69%) have not yet been found outside the region. This number of 'apparent endemics' may decrease over time as the region's biodiversity becomes better known, but it still represents an extraordinarily high level of endemism compared to other sponge genera and other phyla. For example, Van Soest (1994b) found about 24% of all sponges collected from the Seychelles and Amirante islands were endemic to the W Indian Ocean fauna, and Richmond (1997) suggested that only about 15% of all the regional marine biota may be endemic to the W Indian Ocean region.

Biogeographic affinities of the sponge fauna are thought to be essentially Tethyan (Van Soest,

1994a), with southern incursions of species of Gondwanan origin (Hooper & Lévi, 1994; Hooper, 1996a). This fauna is thought to be most similar to the central Indian Ocean and Indo-Malaya regions (Van Soest & Hajdu, 1997), and in this regard follows the general pattern seen in other marine invertebrate phyla (e.g. scleractinarian corals; Veron, 1986; Veron, 1993). Richmond (1997) suggested that about 35% of the West Indian Ocean marine biota was widely distributed, extending into the Red Sea and perhaps as far as the Indo-Malaya region, 35% ranged across the whole Indo-Pacific region, and 15% extended into the warm temperate regions of the Atlantic Ocean. To date only one microcionid, *Clathria (Thalysias) vulpina* (Lamarck), appears to be truly Indo-Pacific 'cosmopolitan', with a confirmed distribution extending from Tonga to the Red Sea (i.e. with conspecificity confirmed through morphological comparisons between recent collections of living populations; Hooper, 1996a). However, even this finding has yet to be confirmed through genetic analysis to determine whether slight morphological differences between regional populations represent intraspecific variability or indicate the existence of a series of possible allopatric sibling species.

Van Soest (1994b) also noted a high level of regional heterogeneity between sponge faunas of the Seychelles and Amirante Island groups, with only 17% of species common to both regional faunas. This supports a similar finding from

TABLE 1. Literature on the sponges of the Western Indian Ocean, Southeast Africa and Arabian Gulf-Red Sea provinces.

Province	Locality	Sponge literature
Coastal East Africa	Zanzibar	Lendenfeld, 1897; Baer, 1905; Jenkin, 1908; Burton, 1959; Thomas, 1976a, 1979b; Pulitzer-Finali, 1993; Kelly, 1997; Magnino & Gaino, 1998
	Tanzania	Jenkin, 1908; Thomas, 1976a; Pulitzer-Finali, 1993; Magnino & Gaino, 1998
	Mozambique	Lévi, 1964; Thomas, 1979a, 1979c, 1980a, 1980b; Laghi et al., 1984; Schmidt et al., 1997
	Kenya	Marsden, 1975; Bruce, 1976; Thomas, 1981a; Vacelet et al., 1991; Pulitzer-Finali, 1993
	Somalia	McCabe et al., 1982; Finamore et al., 1983; Hooper, 1996a
Offshore East Africa	South Africa	Ehlers, 1870; Carter, 1871; Gray, 1873; Vosmaer, 1880; Kirkpatrick, 1900, 1901, 1902a, 1902b, 1903, 1904, 1908, 1913; Sollas, 1908; Stephens, 1915; Burton, 1926, 1929, 1931, 1933a, 1933b, 1936, 1958; Lévi, 1963, 1967; Borojevic, 1967; Day, 1981; Schleyer, 1991; Pettit et al., 1993b; Rudi et al., 1993, 1994a, 1994b, 1995; Barkai et al., 1996; Hooper et al., 1996; Samaai, 1997; Beukes et al., 1998; Koren Goldshlager et al., 1998; McPhail et al., 1998
	Madagascar	Bosraug, 1913; Decary, 1946; Lévi, 1956; Vacelet & Vasseur, 1965, 1966, 1971, 1977; Vacelet, 1967a, 1967b, 1977; Vacelet et al., 1976; Ivanova et al., 1993; Hooper, 1996b
	Aldabra	Lévi, 1961
	Comoros	Sarà et al., 1993b; Pettit et al., 1993a, 1994a, 1994b
	Réunion	Lévi, 1986; Aknin et al., 1996
	Mauritius	Thomson, 1868; Duncan, 1880; Topsent, 1890; Dendy, 1922; Van Soest, 1993
	Saya de Malha	Dendy, 1922; Kolbasov, 1992
	Seychelles	Wright, 1881; Ridley & Dendy, 1887; Topsent, 1893a; Dendy, 1922; Lévi, 1961; Thomas, 1973, 1979c, 1981b; Hooper & Krasochin, 1989; Ngoc Ho, 1990; James et al., 1991; Venkateswarlu et al., 1991; Van Soest, 1994b; Van Soest et al., 1994; Trimurtulu & Faulkner, 1994; Hooper, 1996a; Pettit et al., 1997
	Amirante	Carter, 1880; Ridley, 1884; Ridley & Dendy, 1887; Dendy, 1922; Van Soest et al., 1994; Braekman et al., 1998
Northwest Indian Ocean	Red Sea	Keller, 1889, 1891; Topsent, 1892; Row, 1911; Lévi, 1958, 1965; Burton, 1959; Delseth et al., 1979; Sarà et al., 1979; Kashman et al., 1982, 1989; Mergner, 1982; Sokoloff et al., 1982; Mebs, 1985; Carmely & Kashman, 1986; Vine, 1986; Gebreyesus et al., 1988; Ilan & Loya, 1988, 1990; Carmely et al., 1990; Kolbasov, 1990; Isaacs & Kashman, 1992; Rinkevich et al., 1993; Rudi & Kashman, 1993; Kelli-Bornes & Vacelet, 1995; Guo et al., 1996, 1997a, 1997b; Ramadan, 1997; Beer & Ilan, 1998; Wörheide, 1998
	Ethiopia	Isaacs et al., 1991
	Eritrea	Hooper, 1996a
	Arabian Sea	Carter, 1869; Topsent, 1893b; Dendy, 1913, 1915, 1916a, 1916b, 1916c, 1922; Kumar, 1924a, 1924b, 1924c, 1925; Burton & Rao, 1932; Burton, 1959; Thomas, 1975, 1976b, 1979b, 1988, 1989; Rahim, 1979; Kamat et al., 1981; Patel et al., 1985; Kondracki & Guyot, 1987; James et al., 1989; Parameswaran et al., 1989, 1992a, 1992b, 1994, 1997; Kobayashi et al., 1992a, 1992b; Pettibone, 1993; Sarà & Bavestrello, 1995; Bavestrello et al., 1996; Thomas et al., 1997
	Oman	Sarà & Bavestrello, 1995; Bavestrello et al., 1996

sponge surveys of NW and NE Australian reefs (Hooper, 1994; Hooper et al., 1999), with the implication that taxonomic inventories of regional sponge faunas are largely incomplete, with possibly many new taxa remaining to be discovered within these highly heterogeneous and 'apparent endemic' regional populations. The present study describes one such species discovered during routine surveys of Zanzibar undertaken by MK as part of a project to produce an inventory of the marine fauna and flora and popular field guide to the region (Richmond, 1997).

Methods for preparation and examination of material are described by Hooper (1996a). Spicule measurements refer to (minimum-(mean)-maximum) dimensions of lengths and widths taken from 25 random samples of each spicule category

and are given in micrometres unless otherwise stated. Abbreviations: BMNH, The Natural History Museum, London; QM, Queensland Museum, Brisbane. MK is grateful to Dr Matthew Richmond for facilitating her participation in the East Africa marine surveys.

SYSTEMATICS

PORIFERA Grant
 DEMOSPONGIAE Sollas
 POECILOSCLERIDA Topsent
 MICROCIONINA Hajdu, Van Soest & Hooper
 MICROCIONIDAE Carter

Clathria Schmidt, 1862

Clathria (*Microciona*) Bowerbank, 1862
 Refer to synonymy in Hooper (1996a)

Clathria (Microcionia) richmondi sp. nov.
(Figs 1-4)

ETYMOLOGY. For Dr Matthew D. Richmond, Institute of Marine Sciences, Zanzibar, in recognition of his substantial contribution towards documenting the marine flora and fauna of the E. African coastline (Richmond, 1997).

MATERIAL. HOLOTYPE. QMG306785 (fragment BMNH 1995.6.29.96), E side of Pange Sandbank Reef, Zanzibar Town, Unguja I., Tanzania, 6°10.0'S, 39°9.3'E, 10m depth, 7.viii.1995. coll. M. Kelly. SCUBA.

DISTRIBUTION. Known only from the fringing reefs off Zanzibar Town, Unguja Island, encrusting dead coral substrate on a shallow fringing reef.

DESCRIPTION. Shape. Very thinly encrusting (0.2-0.7mm thick) in small patches (10-20cm diameter) or completely enveloping coral rubble.

Colour. Royal blue with a violet tinge alive, brownish-orange in ethanol.

Oscules. Large (up to 5mm diameter), raised on membranous lip (approximately 4mm high), scattered over entire surface and with prominent, vein-like, radial subsurface drainage canals radiating towards each oscule; oscules and drainage canals collapsed upon preservation.

Texture and Surface Characteristics. Slimy, very smooth, fleshy surface.

Ectosome. No special category of ectosomal spicules present, although choanosomal principal styles arising from the underlying skeleton, standing perpendicular to the substrate, may protrude a long way through the surface. Bundles of toxodragmata occasionally lie on the surface, although most of these appear to be confined below the peripheral skeleton.

Subectosome. Below the surface are plumose bundles of auxiliary subtylostyles, mostly running perpendicular and paratangential to, or occasionally protruding through, the surface. These subectosomal skeletal bundles form stellate brushes associated with (or parallel to) the larger protruding choanosomal principal styles. Toxodragmata form thick bundles below the peripheral skeleton, resembling megasclere spicule tracts, lying tangential to the surface and scattered between the erect choanosomal principal styles.

Choanosome. Microcionid skeletal structure, with thin hymedesmioid basal layer of spongin fibre, approximately 50 thick, highly collagenous, granular, dark brown pigmented, with

calcite detritus embedded beneath; basal spongin with sparsely dispersed echinating acanthostyles embedded and standing perpendicular to substrate, and bulbous spongin fibre nodes up to 150 thick found only in thicker sections of the encrustation; each bulbous fibre node discrete, erect, without any anastomoses between adjacent nodes, and each with 1-5 choanosomal principal styles embedded and perpendicular to substrate, with spicules diverging slightly, becoming plumose towards surface and protruding up to 350 through ectosome; smaller echinating acanthostyles confined mostly to hymedesmioid basal spongin fibre, rarely seen on bulbous fibre nodes; conversely, principal styles only seen on bulbous fibre nodes and not on hymedesmioid basal fibres. Dense horizontal bands of accolada toxodragmata (up to 70 thick) occur about midway through the choanosomal skeleton cross-section, and also in the ectosomal region, lying on or below the surface; few single toxas observed in the mesohyl mostly comprising wing-shaped forms, whereas most toxas forming dragmata. Palmate isochelae moderately abundant within mesohyl; collagen within mesohyl dense, relatively smooth, moderately heavily pigmented orange-brown; choanocyte chambers elongate-oval, up to 40 × 12.

Megascleres. Choanosomal principal styles and subtylostyles very long, slender, slightly curved near basal end, long tapering fusiform points, base either slightly constricted or non-tylote, smooth or very occasionally with anisoxeote terminations. Length 178-(403.7)-622, width 6-(10.2)-12.

Subectosomal auxiliary subtylostyles very long, very slender, straight, fusiform points, with well developed, entirely smooth subtylote bases. Length 198-(351.5)-428, width 2-(3.8)-5.

Echinating acanthostyles relatively uncommon, short, slender, straight or very slightly curved at centre, fusiform-pointed, with moderately well-developed basal constriction; shaft and base with vestigial granular spines confined mainly to basal half of spicule. Length 58-(89.6)-134, width 3-(5.4)-8.

Microscleres. Palmate isochelae moderately common, well-silicified, with thick, well-developed alae comprising over 70% of spicule length. Length 14-(15.2)-17.

Toxas in two forms: Accolada toxas extremely abundant, exceptionally long and very slender, with slight central curvature and straight




FIG. 1. Megascleres from *Clathria (Microcionia) richmondi* sp. nov. (holotype QMG306785). A-B, Choanosomal principal style and subtyle base. C-E, Larger echinating acanthostyle, base and shaft, illustrating mostly smooth, tuberculate ornamentation. F-H, Smaller echinating acanthostyle, base and shaft with rudimentary small spines. I-J, Subectosomal auxiliary subtyle and smooth base.

(non-reflexed) arms, invariably forming toxodragmata. Length 262-(501.3)-975, width 1.5-(1.9)-2.0. Shorter toxas present but uncommon, intermediate between wing-shaped and accolada in morphology, with slight to moderate central curvature, slightly reflexed arms; found in toxodragmata together with accolada toxas and also occasionally singly within the mesohyl. Length 84-(114.8)-154, width 0.8-(1.04)-1.5.

REMARKS. *Clathria (Microcionia) richmondi* sp. nov. is unusual in having 1) huge accolada toxas in dragmata, forming dense bands both within the mesohyl and lying tangential to the surface; 2) a second, less common and much smaller form of toxas, intermediate between wing-shaped and accolada morphology, scattered singly within the mesohyl; 3) a skeleton composed of hymedesmioid basal spongin fibres in thinner sections and microcionid bulbous spongin fibre nodes in thicker parts of the

skeleton, each node with one or few choanosomal principal styles perpendicular to the surface; 4) relatively uncommon echinating acanthostyles, with rudimentary spination, apparently confined to the hymedesmioid basal skeleton; and 5) distinctive field characteristics including a royal blue colour, large oscules with a prominent raised 'lip' and prominent subsurface drainage canals radiating towards each oscule. Although individually these distinctive characters are not unique amongst known species of *Clathria*, in combination they clearly differentiate the Zanzibar species from others.

1) Microsclere morphology, including toxas, appears to be a relatively consistent and useful character to differentiate between similar species (Hooper, 1996a). Six species of *Clathria* have been recorded with accolada toxas forming toxodragmata (*C. (Thalysias) cactiformis* (Lamarck), *C. (Microcionia) densa* (Burton),




FIG. 2. Microscleres from *Clathria (Microcionia) richmondi* sp. nov. (holotype QMG306785). A, Longer accolada toxas. B, Smaller toxas, intermediate between wing-shaped and accolada forms. C, Palmate isochelae.

C. (T.) lendenfeldi Ridley & Dendy, *C. (T.) oxitoxa* Lévi, *C. (Axociella) thetidis* (Hallmann) and *C. (T.) vulpina* (Lamarck) (Hooper, 1996a)); refer to Table 2 for known distributions. In *C. richmondi* the accolada toxas are exceptionally large compared to most species of *Clathria* (262-975 long), and also form very prominent bands within the skeleton. By comparison, those of *C. (T.) cactiformis* are 8-355 long, *C. (M.) densa* (205-305), *C. (T.) lendenfeldi* (7-361) and *C. (T.) vulpina* (8-200), and are scattered throughout the skeleton in loose bundles not forming skeletal tracts. In contrast, accolada toxas of *C. (A.) thetidis* and *C. (T.) oxitoxa* are much larger than those of *C. (M.) richmondi* (175-1280 and 170-3000 long, respectively), and moreover those of *C. (T.) oxitoxa* also form skeletal tracts

within the skeleton (Lévi, 1963). In this regard *C. oxitoxa* is most similar to *C. richmondi*, although all three taxa differ in virtually every other respect (see redescription of *C. oxitoxa* below and *C. thetidis* in Hooper (1996a)).

2) Many species of *Clathria* have two toxa morphologies, and this feature is probably of little diagnostic importance above the species level. Of the species mentioned above only *C. densa* and *C. vulpina* lack both morphologies of toxas.

3) Hymedesmioid – microcionid skeletal structure has been used in the past as a primary diagnostic character for several nominal microcionid genera (e.g. *Axocielita* de Laubenfels, *Hymantho* Burton, *Leptoclathria* Topsent), although this view is no longer widely held (Van

Soest, 1984; Hooper, 1996a). Within *Clathria* these species are now placed in either the subgenera *Microciona* or *Thalysias*, depending on whether ectosomal specialisation is absent or present, respectively. Worldwide there are hundreds of encrusting microcionid species with hymedesmioid and/or microcionid spongin fibre skeletons and a mineral skeleton composed of perpendicular and/or plumose spicule tracts, of which 22 occur in this region (Table 2; species annotated (2)).

4) The presence or absence of echinating spicules, the degree to which they are smooth or spined, and the morphology of spines have been used as generic characters within Microcionidae at one time or another (e.g. *Anaata* de Laubenfels, *Axociella* Hallmann, *Folitispa* de Laubenfels, *Isociella* Hallmann, *Ophlitaspongia* Bowerbank, *Paratenaciella* Vacelet & Vasseur, *Tenaciella* Hallmann). The absence of echinating megascleres remains a valid diagnostic character (at the subgeneric level) for some taxa (e.g. *Isociella*, *Axociella*, *Ophlitaspongia*), by virtue of the consistent combination of this feature and the possession of unusual skeletal structures characterising each of the taxa (Hooper, 1996a; Howson & Chambers, 1999). Within *Clathria* s.s. there are also several species that have lost echinating spicules (e.g. *C. (C.) paucispicula* (Burton), *C. (T.) craspedia* Hooper). By comparison, the absence (loss), rudimentary development and shape of spines on echinating spicules vary widely amongst the many hundreds of species of *Clathria*, although these features appear to be consistent at the species level (e.g. *C. (M.) aceratoobtusata* (Carter) with virtually




FIG. 3. Skeletal structure of *Clathria (Microciona) richmondi* sp. nov. (holotype QMG306785). A, Hymedesmioid-microcionid basal skeleton with tracts of auxiliary subtylostyles lying in multispicular bands within the choanosome. B, Erect echinating acanthostyles (mostly smooth) perpendicular to the hymedesmioid basal skeleton.

smooth spicules, *C. (C.) kylista* Hooper & Lévi with rudimentary spination, *C. (T.) dubia* (Kirkpatrick) with large, heavy spines). Within this continuum *C. richmondi* is most similar to the *C. kylista* condition.

5) As far as can be ascertained from the literature and from personal experience with the Microcionidae of the Indo-Pacific no other species

has a deep royal blue colour in life, but the importance of this character is limited by the lack of good knowledge of their living characters, most taxa known only from preserved specimens. Prominent raised oscules and a subsurface water canal system (radiating towards the oscules and producing a stellate surface pattern) are features common to many encrusting sponges (in which the aquiferous system is marginalised onto the external surface by virtue of the reduced choanosomal thickness), whereas in many microcionids this drainage system often has a different pigmentation from the adjacent ectosome.

Of the 64 species of *Clathria* recorded from the Western Indian Ocean, Southeast Africa and Arabian Gulf-Red Sea provinces 19 have *accolada toxas* (Table 2; species annotated (1)). Most of these species differ substantially from *C. richmondi* in major features such as growth form, skeletal structure, spicule geometries, spicule sizes, possession of specialised ectosomal skeleton (i.e. *Thalysias* condition) etc., whereas in 5 species these differences are more subtle.

Clathria (T.) oxitoxa Lévi (1963) is erect, bushy, flattened lamellate branches, yellow alive; skeleton plumo-reticulate with fibres irregularly cored by very large principal styles and bundles (dragmata) of large oxete toxas scattered throughout the sponge skeleton; ectosomal skeleton with plumose brushes of both ectosomal and subectosomal styles; principal styles regularly curved, with smooth non-tylote bases (450-800 × 35-40); subectosomal auxiliary subtylostyles with microspined bases (350-500 × 9); ectosomal auxiliary subtylostyles similar (100-150 × 4); echinating acanthostyles entirely spined with large spines (75-90 × 10); palmate isochelae in 2 size classes (6 and 13-14); *accolada toxas* nearly oxete, ranging from hair-like and faintly curved arms (170-250 × 0.5-1), thickly oxete with straight arms (750-1300 × 4-7) and extremely long oxetes with straight arms and only slight central curvature (2400-3000 × 10-11); smaller wing-shaped toxas with large central curvature (35-40 × 0.5). This species is most similar to *C. (M.) richmondi* in having exceptionally large *accolada toxas* in dragmata forming skeletal bands, but differs significantly in its live colouration, growth form, skeletal structure and the geometry and size of all spicules (in particular the huge upper size range of *accolada toxas*).

Clathria (M.) densa (Burton, 1959) is massive, dense choanosomal skeleton with semi-plumose

ascending spongin fibres cored by principal styles, interconnected by few transverse fibres and spicule tracts; dense ectosomal skeleton of auxiliary subtylostyles; principal styles with subtylote spined bases (175-298 × 18-35); subectosomal auxiliary subtylostyles with thick spined bases (130-275 × 4-8), echinating acanthostyles thick, slightly curved, heavily spined with aspinose neck (118-156 × 9-16); palmate isochelae (9-13); hair-like *accolada toxas*, distinctly sinuous and raphidiform forming dragmata (205-305 × 0.5-1.5). This species is a borderline case between the subgenera *Microcionia* and *Clathria* given that its choanosomal skeleton is a well-developed microcionid architecture that is verging on reticulate given the existence of vestigial interconnecting fibres and spicule tracts. It also differs from *C. (M.) richmondi* in spicule geometry, spicule sizes, and absence of smaller wing-shaped toxas: in fact the two species are only similar in possessing *accolada toxas* forming dragmata.

Clathria (C.) inhaensis Thomas (1979b) is thinly encrusting, surface conulose; ectosome reduced; choanosomal skeleton reticulate, with well developed ascending primary fibres cored by plumose tracts of principal styles, interconnected by thinner transverse fibres in which few spicules are found and fully embedded within fibres, and both echinated by acanthostyles; subtylostyles interstitial and in brushes arising from tips of main fibres; principal styles with smooth bases (121-172 × 4-5); subectosomal subtylostyles with smooth bases (124-181 × 2-4); echinating acanthostyles with variably spined shaft and spined bases (41-58 × 3-5); palmate isochelae (8-10); *accolada toxas* hair-like (110-145 × 0.5-1.5). This species clearly sits within subgenus *Clathria* given its possession of a reticulate skeletal architecture. It also differs significantly from *C. (M.) richmondi* in spicule geometry, spicule size and absence of toxodragmata.

Clathria (T.) longitoxa (Hentschel, 1912) ranges from thinly encrusting to massive growth form; hymedesmioid to closely reticulate skeleton of stout fibres, with larger and smaller principal styles coring fibres in plumose arrangement, echinated by acanthostyles; subectosomal auxiliary subtylostyles scattered; principal styles curved with smooth non-tylote bases (592-840 × 22-26); smaller principal styles with subtylote granular bases (120-408 × 8-20); subectosomal auxiliary subtylostyles with faintly microspined bases (430-584 × 4-9); ectosomal auxiliary subtylostyles similar (190-320 × 3-5); echinating

acanthostyles evenly spined with spined points ($64-80 \times 6-7$); palmate isochelae ($12-20$); accolada toxas with central U-bend and straight arms ($400-820 \times 1-2$). This species differs from *C. (M.) richmondi* in most respects, showing similarities only in growth form, skeletal structure and possession of accolada toxas.

Clathria (C.) oculata Burton (1933a) has an erect branching anastomosing growth form, drab colouration with tinges of occasional purple; skeletal architecture composed of a subsidiectyal reticulation of spongin fibres fully cored by principal styles and evenly echinating by acanthostyles; principal styles with smooth non-tylote bases ($140-7$); subectosomal auxiliary subtylostyles with smooth bases (160×3); echinating acanthostyles evenly spined with small spines (65×4); accolada toxas slightly curved (160 long); palmate isochelae very small (6 long). This species is only similar to *C. (M.) richmondi* in possessing accolada toxas, differing in most other features.

In addition to these species there are three unnamed species described from Madagascar by Vacelet & Vasseur (1971) showing similarities to *C. richmondi* in the morphology of their accolada toxas, skeletal structure and growth form, although differing in most all other characters.

Clathria (T.) sp. 4 (Vacelet & Vasseur, 1971; see Table 2) is thinly encrusting, yellow alive; choanosomal skeleton microcionid with columns of fibres cored by principal styles and acanthostyles; ectosomal specialisation with some surface brushes but these are not thick; principal styles very slightly subtylote, smooth bases

($130-440 \times 6-12$); subectosomal auxiliary subtylostyles with smooth bases ($150-320 \times 4$); ectosomal auxiliary subtylostyles with microspined bases ($110-200 \times 3$); echinating acanthostyles slightly subtylote, poorly developed spines ($55-60 \times 5$); palmate isochelae in 2 size classes, the smaller contort (5 and 12.5 long); accolada toxas nearly



FIG. 4. Ectosomal skeleton of *Clathria (Microcionia) richmondi* sp. nov. (holotype QMG306785). A, Bundles of subectosomal auxiliary subtylostyles paratangential to and protruding through the surface, loosely associated with erect principal styles. B, Toxodragmata (bundles of accolada toxas) lying on or close to the surface.

TABLE 2. List of *Clathria* species recorded from the Western Indian Ocean, Southeast Africa and Arabian Gulf-Red Sea provinces. Refer to Hooper (Hooper, 1996a) for full synonymy and taxonomic references. Annotation: 1 = species with accolada toxas; 2 = encrusting species with hymedesmioid-microcionid skeletal structure; 3 = identification has yet to be confirmed from examination of voucher specimen; 4 = identification unconfirmed, specimen voucher material missing; 5 = new combination; 6 = currently unrecognisable.

Current taxonomic assignment	Published name	Author	Western Indian Ocean records	Other known distribution
<i>C. (Thalysias) abietina</i> (Lamarck)	<i>C. aculeata</i> Ridley	Burton (1959), Vacelet et al. (1976, 1977)	Red Sea, S Arabian coast, Madagascar	Tropical Australia, central NW Pacific, Philippines
<i>C. (Microcionia) affinis</i> (Carter) ²	<i>M. affinis</i> Carter	Burton (1959)	S Arabian coast, Zanzibar	Gulf of Manaar
<i>C. (Thalysias) amirantiensis</i> Hooper ¹	<i>Colloclathria ramosa</i> Dendy (preocc.)	Dendy (1922), Hooper (1996)	Amirante, Coëtivy, Seychelles	-
<i>C. (Thalysias) anomala</i> (Burton) ¹	<i>R. anomala</i> Burton	Burton (1933)	S South Africa	-
<i>C. (Thalysias) anonyma</i> (Burton) ^{2,5}	<i>M. anonyma</i> Burton	Burton (1959)	Zanzibar	-
<i>C. (Clathria) arbuscula</i> (Row)	<i>Ophlitaspongia arbuscula</i> Row, <i>O. horrida</i> Row	Row (1911)	Red Sea	-
<i>C. (Microcionia) atrasanguinea</i> (Bowerbank) ²	<i>M. atrasanguinea</i> Bowerbank	Carter (1880), Dendy (1922), Burton & Rao (1932), Lévi (1965), Van Soest (1993)	Seychelles, Red Sea, Arabian Sea, Mauritius	Caribbean, NE Atlantic, Mediterranean, coast of India, Gulf of Manaar, Bay of Bengal, Andaman Sea
<i>C. (Clathria) axociona</i> Lévi	<i>C. axociona</i> Lévi	Lévi (1963)	S South Africa	Namibia
<i>C. (Thalysias) cactiformis</i> (Lamarck) ¹	<i>Rhaphidophlus typicus</i> (Carter), <i>C. (T.) cactiformis</i> (Lamarck), <i>Rhaphidophlus</i> sp. 2; Vacelet & Vasseur	Vacelet et al. (1971, 1976, 1977), Hooper (1996)	Madagascar, Somalia, E Africa, Seychelles, Red Sea	S, W & E coasts of Australia
? <i>C. (Clathria) caespes</i> (Ehlers) ⁶	<i>Scopalina caespes</i> (Ehlers)	Hooper (1996)	S South Africa	-
<i>C. (Wilsonella) cercidochela</i> Vacelet & Vasseur	<i>Clathriopsamma cercidochela</i> Vacelet & Vasseur	Vacelet et al. (1971, 1977)	Madagascar	-
<i>C. (Clathria) conica</i> Lévi	<i>C. conica</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Thalysias) cullingworthi</i> Burton	<i>C. cullingworthi</i> Burton	Burton (1931)	Natal	-
<i>C. (Clathria) dayi</i> Lévi	<i>C. dayi</i> Lévi	Lévi (1963)	S South Africa	(? Korea ³)
<i>C. (Thalysias) delaubenfelsi</i> (Lévi)	<i>Rhaphidophlus delaubenfelsi</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Microcionia) densa</i> (Burton) ^{1,2}	<i>M. densa</i> Burton	Burton (1959)	S Arabian coast	-
<i>C. (Clathria) elastica</i> Lévi	<i>C. elastica</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Axociella) fauroti</i> (Topsent)	<i>Axosuberites fauroti</i> Topsent	Topsent (1893)	Gulf of Aden	-
<i>C. (Thalysias) flabellata</i> (Burton)	<i>Rhaphidophlus flabellata</i> Burton	Burton (1936)	S South Africa	-
<i>C. (Clathria) foliascens</i> Vacelet & Vasseur	<i>C. foliascens</i> Vacelet & Vasseur	Vacelet et al. (1971, 1976, 1977)	Madagascar	-
<i>C. (Thalysias) fusterna</i> Hooper	<i>C. fusterna</i> Hooper	Hooper (1996)	Eritrea	N & NE Australia
<i>C. (Clathria) hexagonopora</i> Lévi ¹	<i>C. hexagonopora</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Clathria) indica</i> Dendy	<i>C. indica</i> Dendy	Burton (1931), Thomas (1979)	Natal, Mozambique	SE India, Gulf of Manaar
<i>C. (Clathria) inhacensis</i> Thomas ¹	<i>C. inhacensis</i> Thomas	Thomas (1979)	Mozambique	-
<i>C. (Clathria) irregularis</i> (Burton)	<i>Marleyia irregularis</i> Burton	Burton (1931)	Natal	-

TABLE 2. (cont.)

Current taxonomic assignment	Published name	Author	Western Indian Ocean records	Other known distribution
<i>C. (Clathria) juncea</i> Burton	<i>C. juncea</i> Burton	Burton (1931)	Natal	-
<i>C. (Microciona) laevisissima</i> (Dendy) ²	<i>H. laevisissima</i> Dendy	Dendy (1922)	Mauritius	-
<i>C. (Thalysias) lambda</i> (Lévi)	<i>Leptoclathria lambda</i> Lévi	Lévi (1958)	Red Sea	-
<i>C. (Thalysias) lendenfeldi</i> Ridley & Dendy ¹	<i>C. spicata</i> Hallmann, <i>C. whiteleggii</i> Dendy	Dendy (1922), Burton (1931, 1959), Hooper (1996)	Red Sea, Gulf of Aden, S Arabian coast, Cargados Carajos, Saya de Malha, Somalia, Natal	SE, NE, N. & NW Australia, E Indonesia, Andaman Sea, Gulf of Manaar
<i>C. (Thalysias) lissoclada</i> (Burton)	<i>Rhaphidophlus lissocladius</i> Burton	Lévi (1963)	S South Africa	Falkland Is
<i>C. (Clathria) lobata</i> Vosmaer	<i>C. lobata</i> Vosmaer	Vosmaer (1880), Ridley & Dendy (1887), Stephens (1915), Lévi (1963)	S South Africa	-
<i>C. (Thalysias) longistyla</i> (Burton) ^{2,5}	<i>M. longistyla</i> Burton	Burton (1959)	S Arabian coast	(? Korea ³)
<i>C. (Thalysias) longitoxa</i> (Hentschel) ^{1,2}	<i>M. longitoxa</i> (Hentschel)	Burton (1959)	Gulf of Aden	E Indonesia, Madras
<i>C. (Microciona) microxea</i> (Vacelet & Vasseur) ²	<i>Paratenaciella microxea</i> Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>C. (Wilsonella) mixta</i> Hentschel	<i>C. mixta</i> Hentschel	Burton (1959)	S Arabian coast	E Indonesia ³
<i>C. (Thalysias) nervosa</i> (Lévi)	<i>Axociella nervosa</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Clathria) oculata</i> Burton ¹	<i>C. oculata</i> Burton	Burton (1933, 1959)	Natal	-
<i>C. (Thalysias) oxitoxa</i> Lévi ¹	<i>C. oxitoxa</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Clathria) pachystyla</i> Lévi	<i>C. pachystyla</i> Lévi	Lévi (1963)	S South Africa	-
<i>C. (Axociella) parva</i> Lévi	<i>C. parva</i> Lévi	Lévi (1963)	S South Africa	Namibia
<i>C. (Thalysias) procera</i> (Ridley)	<i>Rhaphidophlus procera</i> Ridley, <i>Echinonema gracilis</i> Ridley	Ridley (1884), Ridley & Dendy (1887), Dendy (1922), Burton & Rao (1932), Burton (1931, 1959), Lévi (1963), Thomas (1973)	Cargados Carajos, Seychelles, Amirante, Red Sea, Arabian coast, Natal	NE, N & NW Australia, E Indonesia, Gulf of Manaar, (? Hawaii ³)
<i>C. (Clathria) raphidotoxa</i> Stephens ¹	<i>C. raphidotoxa</i> Stephens	Stephens (1915), Lévi (1963)	S South Africa	-
<i>C. (Microciona) rhopalophora</i> (Hentschel) ²	<i>M. rhopalophora</i> (Hentschel)	Burton (1959)	Maldives	E. Indonesia, Cocos-Keeling, Gulf of Manaar
<i>C. (Thalysias) robusta</i> (Dendy) ^{1,2}	<i>M. robusta</i> Dendy	Dendy (1922)	Amirante	Singapore
<i>C. (Microciona) seriata</i> (Grant) ^{2,4}	<i>Ophlitaspongia seriata</i> (Grant)	Lévi (1963)	S South Africa	NE. Atlantic, Mediterranean, New Zealand
<i>C. (Clathria) spongodes</i> Dendy	<i>C. spongodes</i> Dendy, <i>C. spongiosa</i> Burton, <i>C. madrepora</i> Dendy	Dendy (1922), Burton (1959), Vacelet et al. (1976)	Red Sea, Gulf of Aden, Amirante, Madagascar, Seychelles	(? Korea ³)
<i>C. (Microciona) stephensae</i> Hooper	<i>M. similis</i> Stephens (preocc.)	Stephens (1915)	S South Africa	-
<i>C. (Microciona) tenuis</i> (Stephens) ²	<i>M. tenuis</i> Stephens	Stephens (1915)	S South Africa	-
<i>C. (Clathria) transiens</i> Hallmann ⁴	<i>C. transiens</i> Hallmann	Burton (1959)	Red Sea	S Australian provinces

TABLE 2. (cont.)

Current taxonomic assignment	Published name	Author	Western Indian Ocean records	Other known distribution
<i>C. (Clathria) typica</i> Kirkpatrick (virtually unrecognisable)	<i>C. typica</i> Kirkpatrick	Kirkpatrick (1904)	Natal	-
<i>C. (Microciona) vacelletia</i> Hooper ²	<i>M. curvichela</i> Vacelet & Vasseur (preocc.)	Vacelet & Vasseur (1965), Hooper (1996)	Madagascar	-
<i>C. (Thalysias) vulpina</i> (Lamarck) ¹	<i>C. frondifera</i> (Bowerbank), <i>C. dichela</i> (Hentschel)	Ridley (1884), Ridley & Dendy (1887), Topsent (1892), Row (1911), Burton (1959), Lévi (1961), Thomas (1973, 1979), Vacelet et al. (1971, 1976, 1977), Pulitzer-Finali (1993), Hooper (1996), Kelly (1997)	Madagascar, Amirante, Seychelles, Red Sea, Mozambique, Aldabra, Zanzibar	Tropical Australia, W & E coasts of India, Gulf of Manaar, Mergui Archipelago, Andaman Sea, Malaysia, E & W Indonesia, N Papua New Guinea, Vietnam, Philippines, Micronesia, S Japan, New Caledonia
<i>C. (Clathria) zoanthifera</i> Lévi	<i>C. zoanthifera</i> Lévi	Lévi (1963)	S South Africa	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur ¹	<i>Rhaphidophlus</i> sp. 1; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur	<i>Rhaphidophlus</i> sp. 3; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur ^{1,2}	<i>Rhaphidophlus</i> sp. 4; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur ^{1,2}	<i>Rhaphidophlus</i> sp. 5; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur ^{1,2}	<i>Rhaphidophlus</i> sp. 6; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Thalysias)</i> sp.; Vacelet & Vasseur	<i>Rhaphidophlus</i> sp. 7; Vacelet & Vasseur	Vacelet et al. (1971, 1977)	Madagascar	-
<i>Clathria (Microciona)</i> sp.; Vacelet & Vasseur ²	<i>Microciona</i> sp. 1; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Microciona)</i> sp.; Vacelet & Vasseur ^{1,2}	<i>Microciona</i> sp. 2; Vacelet & Vasseur	Vacelet & Vasseur (1971)	Madagascar	-
<i>Clathria (Microciona)</i> sp.; Vacelet & Vasseur ²	<i>Microciona</i> sp. 3; Vacelet & Vasseur	Vacelet et al. (1971, 1976)	Madagascar	-
<i>Clathria (Microciona)</i> sp.; Vacelet, Vasseur & Lévi ^{1,2}	<i>Microciona</i> sp. 4; Vacelet, Vasseur & Lévi	Vacelet et al. (1976)	Madagascar	-

oxeote, with slight to virtually no central curvature (85-820 × 0.5-2.5); U-shaped to wing-shaped toxas very small (5-7.5).

Clathria (T.) sp. 5 (Vacelet & Vasseur, 1971) is encrusting, orange alive; hymedesmioid skeleton of principal subtylostyles and acanthostyles erect on basal spongin fibres; subectosomal and ectosomal auxiliary subtylostyles differ only in size and both contribute to both subectosomal and ectosomal surface brushes; principal subtylostyles slender, moderately subtylote microspined bases (140-280 × 5-8); subectosomal auxiliary subtylostyles slightly subtylote, smooth bases (up to 360 × 3); ectosomal auxiliary subtylostyles similar (from 90 × 2); echinating acanthostyles slender, evenly spined (50-60 × 2-3); palmate isochelae in two size classes (4-5 and 12-12.5 long); accolada toxas nearly oxeote,

with straight or only very slightly curved arms and slight angular central flexion (35-250 long).

Clathria (M.) sp. 2 (Vacelet & Vasseur, 1971) is encrusting, pinkish to red alive, with white subectosomal drainage canals clearly visible on the otherwise smooth surface; skeleton microcionid; principal subtylostyles with smooth or microspined bases (330-550 × 13-15); subectosomal auxiliary subtylostyles with smooth bases (120-550 × 2.5-5); echinating acanthostyles slender, entirely spined (100-120 × 5-7.5); palmate isochelae (7.5-20); accolada toxas with only slight curvature of arms and central flexion (130-320); small oxhorn toxas (7.5-20).

LITERATURE CITED

- AKNIN, M., GAYDOU, E.M., BOURYESNAULT, N., COSTANTINO, V., FATTORUSSO, E. & MANGONI, A. 1996. Nor sterols in *Axinella*

- proliferans, sponge from the Indian Ocean. *Comparative Biochemistry and Physiology B Biochemistry & Molecular Biology* 113b(4): 845-848.
- BAER, L. 1905. Silicispongien von Sansibar, Kapstadt und Papeete. *Archiv für Systematische Zoologie* 72:1-32.
- BARKAI, A., DAVIS, C.L. & TUGWELL, S. 1996. Prey selection by the South African Cape rock lobster *Jasus lalandii*: ecological and physiological approaches. *Bulletin of Marine Science* 58(1): 1-8.
- BAVESTRELLO, G., CATTANEO VIETTI, R., CERRANO, C. & SARA, M. 1996. Spicule dissolution in living *Tethya rmanensis* (Porifera: Demospongiae) from a tropical cave. *Bulletin of Marine Science* 58(2): 598-601.
- BEER, S. & ILAN, M. 1998. In situ measurements of photosynthetic irradiance responses of two Red Sea sponges growing under dim light conditions. *Marine Biology* 131(4): 613-617.
- BEUKES, D.R., DAVIES COLEMAN, M.T., KELLY BORGES, M., HARPER, M.K. & FAULKNER, D.J. 1998. Dilemmaones A C, unusual indole alkaloids from a mixed collection of South African sponges. *Journal of Natural Products* 61(5): 699-701.
- BOROJEVIC, R. 1967. Spongiaires d'Afrique du Sud (2) Calcareo. *Transactions of the Royal Society of South Africa* 37: 183-226.
- BOSRAUG, E. 1913. Die Tetractinelliden. *Reise Ostafrika A. Voeltzkow* 3:231-251.
- BOWERBANK, J.S. 1862. On the anatomy and physiology of the Spongiadae. Part III: On the generic characters, the specific characters and the method of examination. *Philosophical Transactions of the Royal Society of London* 152: 1087-1135.
- BRAEKMAN, J.C., DALOZE, D., DE GROOTE, S., FERNANDES, J.B. & SOEST, R.W.M. VAN 1998. New polyketides from the sponge *Plakortis* sp. *Journal of Natural Products* 61(8): 1038-1042.
- BRUCE, A.J. 1976. *Diseias mvitae* sp. nov. a new sponge associate from Kenya (Decapoda, Natantia, Disciadidae). *Crustaceana* 31(2): 119-130.
- BURTON, M. 1926. Descriptions of South African sponges collected in the South African marine survey. Part 1. Myxospongiada and Astrotetrazonida. *Union of South Africa, Fisheries and Marine Biological Survey Report* 4: 1-29.
1929. Description of South African sponges collected in the South African marine survey Part II The Lithistidae, with a critical survey of the desma-forming sponges. *Union of South Africa, Fisheries and Marine Biological Survey Report* (7): 1-12.
1931. On a collection of marine sponges mostly from the Natal coast. *Annals of the Natal Museum* 6(3): 337-358.
- 1933a. Four new marine sponges from Natal. *Annals of the Natal Museum* 7(2): 249-254.
- 1933b. Report on a small collection of sponges from Still Bay, South Africa. *Annals and Magazine of Natural History* (10) 11: 235-244.
1936. Notes on sponges from South Africa, with descriptions of new species. *Annals and Magazine of Natural History* 17: 141-147.
1958. Chapter I. Porifera. *South African Animal Life*. Vol. 5 (Almqvist & Wiksell: Sweden).
1959. Sponges. In *Scientific Reports of the John Murray Expedition 1933-34*. Vol. 10 (British Museum (Natural History): London).
- BURTON, M. & RAO, H.S. 1932. Report on the shallow-water marine sponges in the collection of the Indian Museum. *Records of the Indian Museum* 34(3): 299-356.
- CARMELY, S., GEBREYESUS, T., KASHMAN, Y., SKELTON, B.W., WHITE, A.H. & YOSIEF, T. 1990. Dysidamide, a novel metabolite from a Red Sea sponge *Dysidea herbacea*. *Australian Journal of Chemistry* 43(11): 1881-1888.
- CARMELY, S. & KASHMAN, Y. 1986. Neviotine A, a new triterpene from the Red Sea sponge *Siphonochalina siphonella*. *Journal of Organic Chemistry* 51(6): 784-788.
- CARTER, H.J. 1869. A descriptive account of four subspherical sponges, Arabian and British, with general observations. *Annals and Magazine of Natural History* (4) 4: 1-28.
1871. Description and illustrations of a new species of *Tethya*, with observations on the nomenclature of the Tethyadae. *Annals and Magazine of Natural History* (4) 8: 99-105.
1880. Sponges (including list of sponges dredged by the Birmingham Natural History and Microscopical Society, Falmouth Excursion, 1879. Depth 15-20 Fathoms). *Midland Naturalist* (1880): 55-60, 190-195.
- DAY, J.H. 1981. The estuarine fauna. Pp. 147-178. In *Estuarine ecology with particular reference to southern Africa*. (Balkema: Rotterdam).
- DECARY, R. 1946. Animaux de Madagascar. *Annales du Musée Colonial de Marseille* 6(4): 197-228.
- DELSETH, C., KASHMAN, Y. & DJERASSI, C. 1979. Ergosta-5,7,9(11), 22-tetraen-3B-ol and its 24E-Ethyl Homolog, Two new marine sterols from the Red Sea sponge *Bienina fortis*. *Helvetica Chimica Acta* 62(6): 2037-2045.
- DENDY, A. 1913. Report on the calcareous sponges collected by HMS Sealark in the Indian Ocean. In *Reports of the Percy Sladen Trust Expedition to the Indian Ocean in 1905, Volume 5*. *Transactions of the Linnean Society of London, Zoology* 16: 1-29.
1915. Report on the calcareous sponges collected by Mr James Hornell at Okhamandal in Kattiawar in 1905-1906. In *Report to the Government of Baroda on the Marine Zoology of Okhamandal*. Volume 2. 17: 78-91.
- 1916a. Report on the hexactinellid sponges (triazonida) collected by HMS Sealark in the Indian Ocean. In *Reports of the Percy Sladen*

- Trust Expedition to the Indian Ocean in 1905, Volume 6. Transactions of the Linnean Society of London, Zoology 17: 211-224.
- 1916b. Report on the Homosclerophora and Astrotetaxonida collected by HMS Sealark in the Indian Ocean. In Reports of the Percy Sladen Trust Expedition to the Indian Ocean in 1905, Volume 6. Transactions of the Linnean Society of London, Zoology 17: 225-271.
- 1916c. Report on the non-calcareous sponges collected by Mr James Hornell at Okhamandal in Kattiawar in 1905-1906. In Report to the Government of Baroda on the Marine Zoology of Okhamandal. Volume 2. 17: 96-146.
1922. Report on the Sigmatotetaxonida collected by HMS Sealark in the Indian Ocean. In Reports of the Percy Sladen Trust Expedition to the Indian Ocean in 1905, Volume 7. Transactions of the Linnean Society of London, Zoology 18: 1-164.
- DUNCAN, P.M. 1880. On a parasitic sponge of the order Calcarea. Journal of the Royal Microscopic Society 3:377-383.
- EHLERS, F. 1870. Die Esper'schen Spongien. (Zoologische Sammlung der K. Universität Erlangen. Programm zum Eintritt in der Senat der Königlichen Friedrich-Alexanders-Universität in Erlangen: Erlangen).
- FINAMORE, E., MINALE, L. & ALI MOHAMMED, M. 1983. The stercora of the Somalian sponge *Pterophysilla* (*Pteraplysilla*) *papyracea*. Rendiconto dell'Accademia delle Scienze Fisiche e Matematiche (Serie IV) 50: 81-86.
- GEBREYESUS, T., YOSIEF, T., CARMELY, S. & KASHMAN, Y. 1988. Dysidamide, a novel hexachloro metabolite from a Red Sea sponge *Dysidea* sp. Tetrahedron Letters 29(31): 3863-3864.
- GRAY, J.E. 1873. Natal sponges. Annals and Magazine of Natural History (4) 12: 264.
- GUO, Y., GAVAGNIN, M., MOLLO, E., TRIVEL-LONE, E., CIMINO, G., HAMDY, N.A., FAKHR, I. & PANSINI, M. 1996. A new norsesterterpene peroxide from a Red Sea sponge. Natural Product Letters 9(2): 105-112.
- GUO, Y., GAVAGNIN, M., MOLLO, E. & CIMINO, G. 1997a. Hurghamides A D, new N acyl 2 methylene (beta) alanine methyl esters from Red Sea *Hippospongia* sp. Natural Product Letters 10(2): 143-150.
- GUO, Y., GAVAGNIN, M., MOLLO, E., CIMINO, G., HAMDY, N.A., FAKHR, I. & PANSINI, M. 1997b. Hurghamides A D, new N acyl 2 methylene (beta) alanine methyl esters from Red Sea *Hippospongia* sp. Natural Product Letters 9(4): 281-288.
- HENTSCHEL, E. 1912. Kiesel- und Hornschwämme der Aru und Kei-Inseln. Abhandlungen Senckenbergiana naturforschende Gessellschaft (1912): 295-448.
- HOOPER, G.J., DAVIES-COLEMAN, M.T., KELLY-BORGES, M. & COETZEE, P.S. 1996. New alkaloids from a South African Latrunculid sponge. Tetrahedron Letters 37: 7135-7138.
- HOOPER, J.N.A. 1994. Coral reef sponges of the Sahul Shelf—a case for habitat preservation. Memoirs of the Queensland Museum 36(1): 93-106.
- 1996a. Revision of Microcionidae (Porifera: Poecilosclerida: Demospongiae), with description of Australian species. Memoirs of the Queensland Museum 40: 1-626.
- 1996b. A toxic *Biemna* from Madagascar (Demospongiae: Poecilosclerida). In Willenz, P. (ed.) Recent Advances in Sponge Biodiversity Inventory and Documentation. Bulletin de l'Institut Royal des Sciences Naturelles de Belgique (Supplement) 66: 123-134.
- HOOPER, J.N.A., KENNEDY, J.A., LIST-ARMITAGE, S.E., COOK, S.D. & QUINN, R. 1999. Biodiversity, species composition and distribution of marine sponges in northeast Australia. Pp. 263-274. In Hooper, J.N.A. (ed.) Proceedings of the 5th International Sponge Symposium. Brisbane, June-July 1998. Memoirs of the Queensland Museum 44: 263-274.
- HOOPER, J.N.A. & KRASOCHIN, V.B. 1989. Redescription of the burrowing sponge *Zyzyza massalis* (Dendy) from the Seychelles and Houtman-Abrolhos Islands. The Beagle, Records of the Northern Territory Museum of Arts and Sciences 6(1): 133-140.
- HOOPER, J.N.A. & LÉVI, C. 1994. Biogeography of Indo-west Pacific sponges: Microcionidae, Raspailiidae, Axinellidae. Pp. 191-212. In Soest, R.W.M. van., Kempen, T.M.G. van & Braekman, J.-C. (ed.) Sponges in Time and Space (Balkema: Rotterdam).
- HOWSON, C.M. & CHAMBERS, S.J. 1999. *Ophlitaspongia* and *Ophlitaspongia papilla* reinstated, and a new species of *Ophlitaspongia* described (Porifera: Demospongiae: Microcionidae). Journal of the Marine Biological Association of the United Kingdom 79: 609-620.
- ILAN, M. & LOYA, Y. 1988. Reproduction and settlement of the coral reef sponge *Niphates* sp. (Red Sea). Pp. 745-749. In Choat, J.H., Barnes, D. & Borowitzka, M.A. (eds) Proceedings of the Sixth International Coral Reef Symposium. Townsville, 8-12 August 1988. ICRS, 6th International Coral Reef Symposium Executive Committee 2: 745-749.
1990. Sexual reproduction and settlement of the coral reef sponge *Chalinula* sp from the Red Sea. Marine Biology 105: 25-31.
- ISAACS, S., BERMAN, R., KASHMAN, Y., GEBREYESUS, T. & YOSIEF, T. 1991. New polyhydroxy sterols, dysidamide, and a dideoxyhexose from the sponge *Dysidea herbacea*. Journal of Natural Products, Lloydia 54: 83-91.
- ISAACS, S. & KASHMAN, Y. 1992. Shaagroekol B and C; two hexaprenylhydroquinone disulfates

- from the Red Sea sponge *Toxiclona toxius*. Tetrahedron Letters 33(16): 2227-2230.
- IVANOVA, E.P., MIKHAILOV, V.V., KOUZNETSOVA, T.A., AFIYATULLOV, A.A., KALINOVSKAYA, N.I., ELYAKOV, G.B., KIPRIANOVA, E.A. & GARAGULYA, A.D. 1993. Heterotrophic bacteria associated with the sponge *Dendrilla* sp. and their physiological activity. *Biologiya Morya, Vladivostok* (1993): 3-10.
- JAMES, D.M., KUNZE, H.B. & FAULKNER, D.J. 1991. Two new brominated tyrosine derivatives from the sponge *Druinella* (= *Psammaphysilla*) *purpurea*. *Journal of Natural Products, Lloydia* 54: 1137-1140.
- JAMES, P.S.B.R., GOPINADHA PILLAI, C.S., THOMAS, P.A., JAMES, D.B. & KOYA, S. 1989. 19. Environmental damage and consequences. *Central Marine Fisheries Institute* (1989): 212-227.
- JENKIN, C.F. 1908. The calcareous sponges in the marine fauna of Zanzibar and British East Africa, from collections made by Cyril Crossland, M.A., in the years 1901 and 1902. *Proceedings of the Zoological Society of London* (1908): 434-456.
- KAMAT, S.Y., SOLIMABI, NAQVI, S.W.A., FERNANDES, L., REDDY, C.V.G., BHAKUNI, D.S. & DHAWAN, B.N. 1981. Bioactivity of the extracts from some marine animals of the Indian coast. *Mahasagar* 14(2): 117-122.
- KASHMAN, Y., GROWEISS, A., CARMELY, S., KINAMONI, Z., CZARKIE, D. & ROTEM, M. 1982. Recent research in marine natural products from the Red Sea. *Pure and Applied Chemistry* 54(10): 1995-2010.
- KASHMAN, Y., CARMELY, S., BLASBERGER, D., HIRSCH, S. & GREEN, D. 1989. Marine natural products: new results from Red Sea invertebrates. *Pure and Applied Chemistry* 61(3): 517-520.
- KELLER, C. 1889. Die Spongienfauna des rothen Meeres. I Hälfte. *Zeitschrift für Wissenschaftliche Zoologie* 48: 311-406.
1891. Die Spongienfauna des rothen Meeres. II Hälfte. *Zeitschrift für Wissenschaftliche Zoologie* 52: 294-368.
- KELLY, M. 1997. Porifera. Sponges. Pp. 106-117. In Richmond, M. D. (ed.) *A guide to the seashores of Eastern Africa and the Western Indian Ocean islands* (Sida, Department for Research Cooperation, SAREC. Ord & Vetande AB: Sweden).
- KELLY-BORGES, M. & VACELET, J. 1995. A revision of *Diacarnus* Burton and *Negombata* de Laubenfels (Demospongiae: Latrunculidae) with descriptions of new species from the west central Pacific and the Red Sea. *Memoirs of the Queensland Museum* 38(2): 477-503.
- KIRKPATRICK, R. 1900. Note on the occurrence of the euplectellid sponge *Regadrella phoenix* O. Schmidt, off the South African coast. *Annals of the South African Museum* 13: 63-64.
1901. Description of a new hexactinellid sponge from South Africa. *Annals and Magazine of Natural History* (7) 7: 457-459.
- 1902a. Descriptions of South African sponges. Cape of Good Hope. Department of Agriculture Bulletin. *Marine Investigations in South Africa* 2(14): 219-232.
- 1902b. Descriptions of South African sponges. Part 11. Cape of Good Hope. Department of Agriculture Bulletin. *Marine Investigations in South Africa* 2(14): 171-180.
1903. Descriptions of South African sponges. Part 111. Cape of Good Hope, Department of Agriculture Bulletin. *Marine Investigations in South Africa* 2(16): 233-264.
1904. Sponges. In *Zoological Record (for 1902)*. Abstract by R. von Lendenfeld. *Zoologische Zentralblatt* 10: 147-148.
1908. Description of a new dictyonine sponge from the Indian Ocean. *Records of the Indian Museum* 2: 21-26.
1913. Note on the occurrence of the euplectellid sponge *Regadrella phoenix*, O. Schmidt, off the South African coast. *Annals of the South Africa Museum* 13: 64-66.
- KOBAYASHI, M., CHAVAKULA, R., MURATA, O. & SARMA, N.S. 1992a. Marine terpenes and terpenoids. XIV. Absolute configuration and acid catalyzed transformation of (-)-12, 13-didehydrofurospongins-1 isolated from an Arabian sea sponge, *Fasciospongia cavernosa* Schmidt. *Chemical Pharmaceutical Bulletin (Tokyo)* 40: 599-601.
- KOBAYASHI, M., KRISHNA, M.M., ISHIDA, K. & ANJANEYULU, V. 1992b. Marine sterols. 22. Occurrence of 3 oxo 4,6,8(14) triunsaturated steroids in the sponge *Dysidea herbacea*. *Chemical Pharmaceutical Bulletin (Tokyo)* 40: 72-74.
- KOLBASOV, G.A. 1990. *Acasta pertusa* sp. n. (Cirripedia, Thoracica) from the Red Sea. *Zoologicheskii Zhurnal* 69(9): 142-145.
1992. Two new species of the genus *Acasta* (Cirripedia, Thoracica) from the south western part of the Indian Ocean. *Zoologicheskii Zhurnal* 71(1): 140-145.
- KONDRACKI, M.L. & GUYOT, M. 1987. Smenospongine: a cytotoxic and antimicrobial aminoquinone isolated from *Smenospongia* sp. *Tetrahedron Letters* 28(47): 5815-5818.
- KOREN GOLDSHLAGER, G., KASHMAN, Y. & SCHLEYER, M. 1998. Haliclorsin, a novel diamino alkaloid from the marine sponge *Haliclona tulcarensis*. *Journal of Natural Products* 61(2): 282-284.
- KUMAR, A. 1924a. A new variety of *Leucosolenia gardineri* (Dendy), *Leucosolenia gardineri* var. *vergensis*. *Proceedings of the Lahore Philosophical Society* 3: 21-22.

- 1924b. On a probable new genus of marine sponge from Karachi. Proceedings of the Indian Science Congress 10: 1-167.
- 1924c. Porifera from Karachi. Proceedings of the Lahore Philosophical Society 3: 67-68.
1925. Report on some tetraxonid sponges in the collection of the Indian Museum. Records of the Indian Museum 27: 211-227.
- LAGHI, G.F., MARTINELLI, G. & RUSSO, F. 1984. Localization of minor elements by EDS microanalysis in aragonitic sponges from the St Cassian Beds, Italian Dolomites. *Lethaia* 17(2): 133-138.
- LENDENFELD, R.VON 1897. Spongien von Sansibar. *Abhandlungen Senckenbergiana Naturforschende Gessellschaft* 21: 93-133.
- LÉVI, C. 1956. Spongiaires des côtes de Madagascar. *Mémoires de l'Institut Scientifique de Madagascar (A)* 10: 1-23.
1958. Résultats scientifiques des campagnes de la Calypso'. Fascicule III. V. Campagne 1951-1952 en Mer Rouge. 5. Spongiaires de Mer Rouge recueillis par la Calypso' (1951-1952). *Annales de l'Institut Océanographique, Monaco* 34: 3-46.
1961. Résultats scientifique des campagnes de la Calypso'. Fascicule V. XIV. Campagne 1954 dans l'Océan Indien. 2. Les spongiaires de l'île Aldabra Campagne Océanographique de la Calypso' (May-Juin 1954). *Annales de l'Institut Océanographique, Monaco* 39: 1-31.
1963. Spongiaires d'Afrique du Sud (1) Poccilosclerides. *Transactions of the Royal Society of South Africa* 37(1): 1-72.
1964. Spongiaires du canal de Mozambique. *Bulletin du Muséum National d'Histoire Naturelle* 36(3): 384-395.
1965. Spongiaires recoltés par l'Expedition Israelienne dans le sud de la Mer Rouge en 1962. *Bulletin of the Sea Fisheries Research Station, Israel. Israel South Red Sea Expedition, 1962, Report* 13): 3-27.
1967. Spongiaires d'Afrique du Sud (3) Tétractinellides. *Transactions of the Royal Society of South Africa* 37(3): 227-256.
1986. *Laocaetis perion* nov. sp., spongiaire hexactinellide Craticulariidae de l'océan Indien. *Bulletin du Muséum National d'Histoire Naturelle, Paris* (4) 8(A, 3): 437-442.
- MAGNINO, G. & GAINO, E. 1998. *Haplosyllis spongicola* (Grube) (Polychaeta, Syllidae) associated with two species of sponges from East Africa (Tanzania, Indian Ocean). *Marine Ecology* 19(2): 77-87.
- MARSDEN, J.R. 1975. Classes of lipids in marine sponges from Kenya. *Journal of Experimental Marine Biology and Ecology* 19: 9-18.
- McCABE, T., CLARDY, J., MINALE, L., PIZZA, C., ZOLLO, F. & RICCIO, R. 1982. A triterpenoid pigment with the isomalabaricane skeleton from the marine sponge *Stelletta* sp. *Tetrahedron Letters* 23(33): 3307-3310.
- McPHAIL, K., DAVIES COLEMAN, M.T. & COETZEE, P. 1998. A new furanosesterterpene from the South African nudibranch *Hypselodoris capensis* and a Dictyoceratida sponge. *Journal of Natural Products* 61(7): 961-964.
- MEBS, D. 1985. Chemical defense of a dorid nudibranch, *Glossodoris quadricolor* from the Red Sea. *Journal of Chemical Ecology* 11(6): 713-716.
- MERGNER, H. 1982. Man made influences on and natural changes in the settlement of the Aqaba reefs (Red Sea). *Proceedings of the International Coral Reef Symposium* (4): 193-207.
- NGOC HO, N. 1990. Nine Indo Pacific species of *Upogebia* Leach (Crustacea: Thalassinidea: Upogebiidae). *Journal of Natural History* 24(4): 965-985.
- PARAMESWARAN, P.S., DAS, B. & KAMAT, S.Y. 1994. Lipid contents of the sponge *Haliclona* sp. *Indian Journal of Chemistry (B) Organic Chemistry, Medicinal Chemistry* 33B(1): 99-101.
- PARAMESWARAN, P.S., KAMAT, S.Y., CHANDRAMOHAN, D., NAIR, S. & DAS, B. 1992a. Anti-bacterial compounds from the sponge *Haliclona* sp. Pp. 417-420. In Desai, B.N. (ed.) *Oceanography of the Indian Ocean* (Oxford & IBH Publishing Co.: New Delhi, Bombay).
- PARAMESWARAN, P.S., NAIK, C.G., DAS, B. & KAMAT, S.Y. 1989. Sterols from the Lakshadweep sponge, *Ircinia ramosa* (Keller). *Indian Journal of Chemistry Section (B) Organic Chemistry, Medicinal Chemistry* 28(12): 1091-1092.
- 1992b. Minor sterols from the sponge *Ircinia ramosa* (Keller). Pp. 413-416. In Desai, B.N. (ed.) *Oceanography of the Indian Ocean* (Oxford & IBH Publishing Co.: New Delhi, Bombay).
- PARAMESWARAN, P.S., NAIK, C.G. & HEGDE, V.R. 1997. Secondary metabolites from the sponge *Tedania anhelans*: isolation and characterization of two novel pyrazole acids and other metabolites. *Journal of Natural Products, Lloydia* 60(8): 802-803.
- PATEL, B., PATEL, S. & BALANI, M.C. 1985. Can a sponge fractionate isotopes? *Proceedings of the Royal Society of London (B) Biological Sciences* 224(1234): 23-41.
- PETTIBONE, M.H. 1993. Scaled polychaetes (Polynoidae) associated with ophiuroids and other invertebrates and review of species referred to *Malmgrenia* McIntosh and replaced by *Malmgreniella* Hartman, with descriptions of new taxa. *Smithsonian Contributions to Zoology* (538): 1-92.
- PETTIT, G.R., CICHACZ, Z.A., RUI, T., HOARD, M.S., MELODY, N. & PETTIT, R.K. 1998a. Antineoplastic agents. 386. Isolation of sesterstains 13 from the marine sponge *Hyrtios erecta*. *Journal of Natural Products* 61(1): 13-16.
- PETTIT, G.R., GAO, F., DOUBEK, D.L., BOYD, M.R., HAMEL, E., BAI, R.L., SCHMIDT, J.M.,

- TACKETT, L.P. & RÜTZLER, K. 1993a. Anticoplastic agents. 252. Isolation and structure of Halistatin-2 from the Comoros Marine Sponge *Axinella carteri*. *Gazzetti Chimica Italiana* 123: 371-377.
- PETTIT, G.R., HERALD, C.L., CICHACZ, Z.A., GAO, F., SCHMIDT, J.M., BOYD, M.R., CHRISTIE, N.D. & BOETTNER, F.E. 1993b. Isolation & structure of the powerful human cancer cell growth inhibitors spongistatins 4 & 5 from an African *Spirastrella spinispirulifera* (Porifera). *Journal of the Chemical Society, Chemical Communications* 1993: 1805-1807.
- PETTIT, G.R., GAO, F., SCHMIDT, J.M., CHAPUIS, J.-C. & CERNY, R.L. 1994a. Isolation and structure of Axinastatin 5 from a Republic of Comoros marine sponge. *Bioorganic and Medicinal Chemistry Letters* 4(24): 2935-2940.
- PETTIT, G.R., RUI, T., HERALD, D.L., CERNY, R.L. & WILLIAMS, M.D. 1994b. Antineoplastic agents. 277. Isolation and structure of phakellistatin 3 and isophakellistatin 3 from a Republic of Comoros marine sponge. *Journal of Organic Chemistry* 59(7): 1593-1595.
- PETTIT, G.R., McNULTY, J., HERALD, D.L., DOUBEK, D.L., CHAPUIS, J.C., SCHMIDT, J.M., TACKETT, L.P. & BOYD, M.R. 1997. Antineoplastic agents. 362. Isolation and X ray crystal structure of dibromophakellistatin from the Indian Ocean sponge *Phakellia mauritiana*. *Journal of Natural Products, Lloydia* 60(2): 180-183.
- PETTIT, G.R., TAN, R., MELODY, N., CICHACZ, Z.A., HERALD, D.L., HOARD, M.S., PETTIT, R.K. & CHAPUIS, J.C. 1998b. Antineoplastic agents. 397. Isolation and structure of sesterstatins 4 and 5 from *Hyrtios erecta* (The Republic of Maldives). *Bioorganic Medical and Chemical Letters* 8: 2093-2098.
- PULITZER-FINALI, G. 1993. A collection of marine sponges from East Africa. *Annali del Museo Civico di Storia Naturale G. Doria* 89: 247-350.
- RAHIM, M.A. 1979. Porifera. *Publications Saudi Biological Society* 3: 46.
- RAMADAN, S.A. 1997. Two new species of mesostigmatid mites (Acari) associated with sponges from the Red Sea, Egypt. *Assiut Veterinary Medical Journal* 38(75): 191-204.
- RICHMOND, M.D. 1997. A guide to the seashores of Eastern Africa and the Western Indian Ocean islands. (Sida, Department for Research Cooperation, SAREC. Ord & Vetandc AB: Sweden).
- RIDLEY, S.O. 1884. Spongiida. Pp. 366-482, 582-635. In *Report on the Zoological Collections made in the Indo-Pacific Ocean during the Voyage of H.M.S. Alert 1881-2* (British Museum (Natural History): London).
- RIDLEY, S.O. & DENDY, A. 1887. Report on the Monaxonida collected by HMS Challenger' during the Years 1873-76. Pp. 1-275. In *Report on the Scientific Results of the Voyage of H.M.S. Challenger during the Years 1873-76*. 20 (Her Majesty's Stationery Office: London, Edinburgh, Dublin).
- RINKEVICH, B., SHASHAR, N. & LIBERMAN, T. 1993. Nontransitive xenogeneic interactions between four common Red Sea sessile invertebrates. Pp. 833-839. In *Richmond, R.H. Proceedings of the Seventh International Coral Reef Symposium*. Mangilao, Guam, 22-26 June 1992. University of Guam Press 2: 833-839.
- ROW, R.W.H. 1911. Reports on the marine biology of the Sudanese Red Sea, from collections made by Cyril Crossland, MA, BSc, FZS. XIX. Report on the sponges collected by Mr Cyril Crossland in 1904-5. Part II. Non-Calcareous. *Journal of the Linnean Society, Zoology* 31(208): 287-400.
- RUDI, A. & KASHMAN, Y. 1993. Aaptosine a new cytotoxic 5,8 diazabenz(c)azulene alkaloid from the Red Sea sponge *Aaptos aaptos*. *Tetrahedron Letters* 34(29): 4683-4684.
- RUDI, A., TALPIR, R., KASHMAN, Y., BENAYAHU, Y. & SCHLEYER, M. 1993. Four new C16 1,2 dioxene polyketides from the sponge *Plakortis aff. simplex*. *Journal of Natural Products, Lloydia* 56(12): 2178-2182.
- RUDI, A., KASHMAN, Y., BENAYAHU, Y. & SCHLEYER, M. 1994a. Sodwanones A-F, new triterpenoids from the marine sponge *Axinella weltneri*. *Journal of Natural Products, Lloydia* 57(10): 1416-1423.
- RUDI, A., STEIN, Z., GREEN, S., GOLDBERG, I., KASHMAN, Y., BENAYAHU, Y. & SCHLEYER, M. 1994b. Phorbazoles A-D, novel chlorinated phenylpyrrolyloxazoles from the marine sponge *Phorbos aff. elathrata*. *Tetrahedron Letters* 35(16): 2448, 2589-2592.
- RUDI, A., KASHMAN, Y., BENAYAHU, Y. & SCHLEYER, M. 1995. Durbinal A, B and C: three new cytotoxic sponge metabolites. *Tetrahedron Letters* 36(27): 4853-4856.
- SAMAAL, T. 1997. Systematics, phylogeny and biogeography of a selection of poecilosclerid sponges from Ouderkraal, on the west coast of South Africa. Unpubl. MSc thesis, Department of Biology, Imperial College, London.
- SARÀ, M. & BAVESTRELLO, G. 1995. *Tethya omanensis*, a remarkable new species from an Oman cave (Porifera, Demospongiae). *Bollettino di Zoologia* 62(1): 23-27.
- SARÀ, M. & CORRIERO, G. 1994. A new species of *Tethya*, *T. tenuisclera* (Porifera Demospongiae) from the Maldives. *Bollettino dei Musei e degli Istituti Biologici della Università di Genova* 58-59: 69-75.
- SARÀ, M., CORRIERO, G. & BAVESTRELLO, G. 1993a. *Tethya* (Porifera, Demospongiae) species coexisting in a Maldivian coral reef lagoon: taxonomical, genetic and ecological data. *Marine Ecology* 14(4): 341-355.
- 1993b. *Tethya peracuta* (Topsent) and *T. comorensis* n. sp. (Porifera, Demospongiae) from the coral

- reef of Mayotte (Comores). *Bollettino di Zoologia* 60: 219-224.
- SARÀ, M., PANSINI, M. & PRONZATO, R. 1979. Zonation of photophilous sponges related to water movement in reef biotypes of Obhor Creek (Red Sea). In Lévi, C. & Boury-Esnault, N. (eds) *Biologie des Spongiaires*. *Sponge Biology. Colloques Internationaux du Centre National de la Recherche Scientifique* (291): 282-288.
- SCHLEYER, M.H. 1991. Shell borers in the oyster, *Striostrea margaritacea*: pests or symbionts? *Symbiosis* 10(1-3): 135-144.
- SCHMIDT, E.O. 1862. *Die Spongien des Adriatischen Meeres* (Wilhelm Engelmann: Leipzig).
- SCHMIDT, E.W., HARPER, M.K. & FAULKNER, D.J. 1997. Mozamides A and B, cyclic peptides from a theonellid sponge from Mozambique. *Journal of Natural Products, Lloydia* 60(8): 779-782.
- SOEST, R.W.M. van 1984. Marine sponges from Curaçao and other Caribbean localities. Part III. Poecilosclerida. *Studies on the Fauna of Curaçao and other Caribbean Islands* (199): 1-167.
1993. Distribution of sponges on the Mauritanian continental shelf. In Wolff, W.J., van der Land, J., Nienhuis, P.H. & de Wilde, P.A.W.J. (eds) *Ecological studies in the coastal waters of Mauritania*. *Hydrobiologia*: 258: 95-106.
- 1994a. Demosponge distribution patterns. Pp. 213-224. In Soest, R.W.M. van, Kempen, T.M.G. van & Braekman, J.-C. (eds) *Sponges in time and space*. (Balkema: Rotterdam).
- 1994b. Sponges of the Seychelles. Pp. 65-74. In van der Land, J. (ed.) *Oceanic reefs of the Seychelles: report on a cruise of RV Tyro to the Seychelles in 1992 and 1993* (National Museum of Natural History: Leiden).
- SOEST, R.W.M. VAN & HAJDU, E. 1997. Marine area relationships from twenty sponge phylogenies. A comparison of methods and coding strategies. *Cladistics* 13: 1-20.
- SOEST, R.W.M. VAN, ZEA, S. & KIELMAN, M. 1994. New species of *Zyzya*, *Cornulella* and *Damiria* (Porifera: Poecilosclerida), with a review of fistular genera of Iophonidae. *Bijdragen tot de Dierkunde* 64(3): 163-192.
- SOKOLOFF, S., HALEVY, S., USIELI, V., COLORNI, A. & SAREL, S. 1982. Pranicin A and B, nor scsterterpenoid peroxide antibiotics from Red Sea sponges. *Experientia, Basel* 38: 337-338.
- SOLLAS, I.B.J. 1908. The inclusion of foreign bodies by sponges, with a description of a new genus and species of Monaxonida. *Annals and Magazine of Natural History* (8) 1: 395-401.
- STEPHENS, J. 1915. Atlantic sponges collected by the Scottish National Antarctic Expedition. *Transactions of the Royal Society of Edinburgh* 50: 423-467.
- THOMAS, P.A. 1973. Marine Demospongiae of Mahe Island in the Seychelles Bank (Indian Ocean). *Annales du Musée Royal de l'Afrique Central-Tervuren, Belgique* (8, Sciences Zoologiques) (203): 1-91.
1975. Boring sponges of Zuari and Mandovi Estuaries. *Bulletin of the Department of Marine Science University of Cochin* 7(1): 117-126.
- 1976a. Marine Demospongiae of Zanzibar Island. *Journal of the Marine Biology Association of India* 18(3): 448-460.
- 1976b. The history of spongology of the Indian Ocean. *Journal of the Marine Biological Association of India* 18(3): 610-625.
- 1979a. Demospongiae of Minicoy Island (Indian Ocean) Part 1 - orders Keratosida and Haplosclerida. *Journal of the Marine Biological Association of India* 21(1-2): 10-16.
- 1979b. *Endectyon lamellosa* n. sp., (Demospongiae: Poecilosclerida, Raspailiidae) from the Indian seas and a revised key to the Indian species of *Endectyon* Topsent. *Journal of the Marine Biological Association of India* 18(1): 169-172.
- 1979c. Studies of sponges of Mozambique Channel 1. Sponges of Inhaca Island. 2. Sponges of Mambone and Paradise Islands. *Annales du Musée Royal de l'Afrique Centrale, Tervuren, Belgique* (8, Sciences Zoologiques). *Koninklijk Museum Voor Midden-Afrika Tervuren Belgie Annalen Zoologische Wetenschappen* (227): 1-73.
- 1980a. Demospongiae of Minicoy Island (Indian Ocean) Part 2 - order Poecilosclerida. *Journal of the Marine Biological Association of India* 22(1-2): 1-7.
- 1980b. Demospongiae of Minicoy Island (Indian Ocean) Part 3 - orders Halichondrida, Hadromerida, Epipolasida and Choristida. *Journal of the Marine Biological Association of India* 22(1-2): 8-20.
- 1981a. Marine Demospongiae of Ras Iwatine (Kenya). *Journal of the Marine Biological Association of India* 18(3): 642-649.
- 1981b. A second collection of marine Demospongiae from Mahe Island in the Seychelles Bank (Indian Ocean). *Koninklijk Museum Voor Midden-Afrika Tervuren Belgie Annalen Zoologische Wetenschappen* (233): 1-54.
1988. Sponge generated bioerosion in Lakshadweep. *Indian Council of Agricultural Research Marine Fisheries Information Service Technical and Extension Series* (86): 20-26.
1989. 13. Sponge fauna of Lakshadweep. *Central Marine Fisheries Research Institute* (1989): 150-161.
- THOMAS, P.A., GOPINADHA PILLAI, C.S. & RAJAGOPALAN, M.S. 1997. Demospongiae of the Gulf of Kutch. *Journal of the Marine Biological Association of India* 38(1-2): 124-132.
- THOMSON, W. 1868. On the vitreous' sponges. *Annals and Magazine of Natural History* (4) 1: 114-132.

- TOPSENT, E. 1890. Etudes de Spongiaires. II. Description d'une Lithistide molle de la Réunion (*Kaliapsis permollis* nov. sp.). Revue de Biologie du Nord France 2(8): 294-296.
1892. Éponges de la Mer Rouge. Mémoires de la Société Zoologique de France 5: 21-29.
- 1893a. Mission Scientifique de M. Ch. Alluaud aux Iles Séchelles (Mars-Mai 1892), Spongiaires. Bulletin de la Société Zoologique de France 18:172-175.
- 1893b. Note sur quelques Éponges du Golfe de Tadjoura recueillies par M le Dr L. Faurot. Bulletin de Société zoologique de France 18: 177-182.
- TRIMURTULU, G. & FAULKNER, D.J. 1994. Six new diterpene isonitriles from the sponge *Acanthella cavernosa*. Journal of Natural Products, Lloydia 57(4): 501-506.
- VACELET, J. 1967a. Descriptions d'éponges Pharétronides actuelles des tunnels obscurs sous-récifaux de Tuléar (Madagascar). Recueil des Travaux de la Station Marine d'Endoume-Marseille (6): 37-62.
- 1967b. Quelques éponges Pharétronides et Silico-Calcaires de grottes sous-marines obscures. Recueil des Travaux de la Station Marine d'Endoume 58: 121-132.
1977. Éponges Pharétronides actuelles et scléro-sponges de Polynésie française, de Madagascar et de La Réunion. Bulletin du Muséum National d'Histoire Naturelle (3, Zoologie) 307(444): 345-368.
- VACELET, J., TIERCELIN, J.J. & GASSE, F. 1991. The sponge *Dosilia browni* (Spongillidae) in Lake Baringo, Gregory Rift, Kenya. Hydrobiologia 211(1): 11-18.
- VACELET, J. & VASSEUR, P. 1965. Spongiaires des grottes et surplombs des récifs de Tuléar (Madagascar). Recueil des Travaux de la Station Marine d'Endoume-Marseille, Supplément (4): 71-123.
1966. Les tunnels obscurs sous-récifaux de Tuléar (Madagascar) et leur faune de spongiaires. Pp. 378. In Abstracts of the Second International Oceanographic Congress. (Moscow).
1971. Éponges des récifs coralliens de Tuléar (Madagascar). Tethys, Supplément 1: 51-126.
1977. Sponge distribution in coral reefs and related areas in the vicinity of Tuléar (Madagascar). Pp. 113-117. In Proceedings of the 3rd International Coral Reef Symposium. Miami, Florida. (Rosenstiel School of Marine and Atmospheric Science: University of Miami).
- VACELET, J., VASSEUR, P. & LÉVI, C. 1976. Spongiaires de la pente externe des récifs coralliens de Tuléar (sud-Ouest de Madagascar). Mémoires du Muséum National d'Histoire Naturelle, Paris (A, Zoologie) 49: 1-116.
- VENKATESWARLU, Y., FAULKNER, D.J., RIOS STEINER, J.L., CORCORAN, E. & CLARDY, J. 1991. Smenochromenes, unusual macrocyclic sesquiterpene hydroquinone derivatives from a Seychelles sponge of the genus *Smenospongia*. Journal of Organic Chemistry 56(22): 6271-6274.
- VERON, J.E.N. 1986. Corals of Australia and the Indo-Pacific. (Angus & Robertson: Sydney, London).
1993. A biogeographic database of hermatypic corals. Species of the Central Indo-Pacific Genera of the world. Australian Institute of Marine Science Monograph Series 10: 1-433.
- VINE, P. 1986. Red Sea invertebrates. (IMMEL Publishing: London).
- VOSMAER, G.C.J. 1880. The sponges of the Leyden Museum. I. The family of the Desmacidinae. Notes from the Leyden Museum 2: 99-164.
- WÖRHEIDE, G. 1998. The reef cave dwelling ultraconservative coralline demosponge *Astrosclera willeyana* Lister 1900 from the Indo-Pacific. Facies 38: 1-88.
- WRIGHT, E.P. 1881. On a new genus and species of sponge (*Alema seychellensis*) with supposed heteromorphic zooids. Transactions of the Irish Academy 28: 13-20.